

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

459 May 4, 1906

Subports.

Mayaguez: 8 vessels inspected and 15 bills of health issued. Arecibo: 2 vessels inspected and 5 bills of health issued. Humacao: No vessels inspected and 4 bills of health issued. Aguadilla: 2 vessels inspected and 9 bills of health issued. Arroyo: 1 vessel inspected and 8 bills of health issued. Fajardo: 10 vessels inspected and 10 bills of health issued.

MORTALITY.

Mayaguez, 124 deaths, 3 of which were due to anemia, 6 to acute bronchitis, 19 to tuberculosis, 11 to uncinariasis, 2 to pneumonia, 5 to intestinal tuberculosis, and 1 to enteric fever.

Arecibo, 83 deaths, 12 due to uncinariasis, 6 to anemia, 10 to infantile tetanus, 13 to tuberculosis, 1 to dysentery, and 1 to pneumonia.

Aguadilla, 39 deaths, 3 due to infantile tetanus, 3 to broncho-pneumonia, 4 to uncinariasis, 6 to tuberculosis, 1 to paludism, 1 to diphtheria, and 1 to meningitis.

Fajardo, 54 deaths, 3 due to paludic fever, 4 to broncho-pneumonia, 2 to enteric fever, 6 to tuberculosis, and 1 to infantile tetanus.

Arroyo, 17 deaths, 1 due to uncinariasis, 5 to paludism, 2 to infantile tetanus, and 3 to pneumonia.

Humacao, no report received.

No quarantinable diseases were reported from the subports.

Mortality at San Juan, March, 1906.

CAUSES OF DEATHS.

Tuberculosis	13	Typhoid-pneumonia	1
Tetanus, infantile	5	Anemia	1
Broncho-pneumonia	2	Gastritis	1
Intestinal tuberculosis	2	Gastro-enteritis	4
Meningitis	5	Entero-colitis	10
Bronchitis, acute		Total from all other causes	71
Dysentery	1		
Pneumonia			120

PORTUGAL.

Report from Lisbon—Smallpox in the Cape Verde Islands.

Vice-Consul Kinchant reports, March 29, as follows:

The consular agent at St. Vincent, Cape Verde Islands, reports that smallpox has been very prevalent for some months in St. Nicholas, and that several immigrants are leaving that island for Brava to take passage for New Bedford, Mass.

He has been directed to be careful in filling out bills of health for vessels clearing from the ports in those islands.

FOREIGN AND INSULAR STATISTICAL REPORTS OF COUNTRIES AND CITIES—UNTABULATED.

Brazil—Pernambuco.—Two weeks ended March 15, 1906. Estimated population, 200,000. Total number of deaths, 306, including enteric fever 3, whooping cough 1, smallpox 11, and 52 from tuberculosis.